



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for SITAPUR District
Chander Shekhar Azad University of Agriculture & Technology
Kanpur, Uttar Pradesh



Agromet Advisory Bulletin

Date : 2025-04-01

Weather Forecast of District SITAPUR (Uttar Pradesh) Issued On : 2025-04-01 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-04-02	2025-04-03	2025-04-04	2025-04-05	2025-04-06
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	38.0	38.0	39.0	40.0	40.0
Tmin(°C)	17.0	18.0	19.0	19.0	19.0
RH-I(%)	40	39	41	40	45
RH-II(%)	18	19	24	25	27
Wind Speed(kmph)	9	6	6	8	8
Wind Direction(Degree)	312	328	337	301	306
Cloud Cover(Octa)	0	3	2	0	0
Warning	No Warning	No Warning	No Warning	No Warning	No Warning

Forecast Summary:

According to the weather forecast received from the Indian Meteorological Department, due to light to moderate cloud cover in the second, third day and clear weather appearing in remaining day, there is a no possibility of rain . The maximum temperature is between 38.0- 40.0°C, which is likely to be 2-3°C higher normal and the minimum temperature is between 17.0-19.0°C, which is likely to be 1-2°C higher normal. The maximum and minimum range of relative humidity is between 39-41 and 18-27%. Wind direction is north-west and wind speed is between 6.0-9.0 kmph with gusts expected to be 2-3 kmph high speed than normal.

Weather Warnings (Valid Till 08:30 IST of the next day)

According to the weather forecast received from the Indian Meteorological Department, there is no warning in next five days.

Likely impacts of weather warnings on Agriculture and associated Agromet advisories

Keeping in view the weather conditions, maintain proper moisture in summer vegetables and crops by doing light irrigation as per requirement. And harvest and thresh mature crops like mustard, gram, peas should be harvested in the morning/evening and do not go out of the house unnecessarily during the afternoon. While working in the field, tie a towel on the head and keep a water bottle with you.

General Advisory:

Farmers are advised to maintain proper moisture in standing vegetables and crops by doing light irrigation as per requirement and harvesting and threshing of mature crops like mustard, gram, peas

should be done in the morning / evening and do not go out of the house unnecessarily during the afternoon. While working in the field, tie a towel on the head and keep a water bottle with you. Keep the grains of threshed crops in a safe place. To prevent infection of thrips insect and purple spot disease in onion, it is advised to keep a regular watch on the crops. Use insecticides, pesticides and weedicides separately or wash the equipment with clean water and do not spray insecticides, pesticides and weedicides while standing in the opposite direction of the wind. If possible, spray in the evening. After spraying, wash hands thoroughly with soap or hand wash before eating or drinking and wash clothes and take a bath.

SMS Advisory:

Farmers are advised to maintain proper moisture in standing vegetables and crops by doing light irrigation as per requirement. Working in the field, tie a towel on the head and keep a water bottle.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
WHEAT	Wheat crop is in the stage of grain filling/dough, which is sensitive to lack of moisture; hence farmers should maintain proper moisture by doing light irrigation as per need. Use fodder made of zinc phosphide or aluminum phosphide cake when rats are found in the wheat crop. Make collective efforts for the prevention of rats.
MAIZE	In the timely sown maize crop, first irrigation should be done 20-25 days after sowing, and weeding should be done when the leaves appear. For controlling the appearance of the larvae of the cabbage borer fly, spray Oxydemeton-methyl 25 EC or Chlorantraniliprole 375 ml. per hectare by dissolving it in 500-700 liters of water.
BLACK GRAM	In the timely sown urad crop, first irrigation should be done 30-35 days after sowing, and weeding should be done when the leaves appear. If infestation of green hopper/thript insect is seen in urad crop, spray Imidacloprid 17.8% SL 2.5 ml per hectare by dissolving it in 500-700 liters of water.
GREEN GRAM	Recommended varieties of moong- Narendra moong-1, Malviya Jagranti, Samrat, moong Janpriya, Meha, HUM-16, Malviya Jyoti, TMV-37, Malviya Jan Chetna, IPM-2-3, IPM-2-14, Kanika, Azad moong-1 Hira, Surya, etc. Arrange for fertilizer and seeds for sowing any one of these varieties and then start sowing.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
ONION	If purple spot disease is seen in onion crop, then for its control, prepare a solution of Mancozeb 0.25% (0.25 gm/liter water) and spray it 3-4 times at an interval of 15-20 days. Complete the sowing of summer vegetables like ladyfinger, zucchini, cucumber, bitter gourd, bottle gourd, pumpkin etc. soon. In case of infestation of red pumpkin insect in pumpkin category vegetables, spray 5% Malathion dust in the soil around the stem of the plants in the morning and spray it on the plants. Prepare the field for transplanting summer crops like tomato, chilli etc. and also transplant the prepared seedlings.
MANGO	Maintain proper moisture by irrigating the mango orchards. There is a possibility of midge insect infestation in mango blossoms. So, to control this, spray by dissolving 1.0 ml of Fenitrothion in one litre of water. There is a possibility of larva and lassi insect infestation in mango orchards. So, to control this, spray by dissolving 2.0 ml of Chlorpyrifos 20 EC or 3.0 ml of Fenitrothion 50 EC in one litre of water.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	After the animals are fed, make 750 grams mustard oil, 100 grams turmeric, 50 grams dry ginger, 50 grams cumin, 500 grams jaggery mixture and OT for morning and evening for three days. In view of the current weather, farmers are advised to animals in the open during the night and draw and put curtains of jut sack on windows and doors at night and remove put curtains in the sun during the day. Give sufficient quantity of grains to the animals with green and dry fodder. Animals must be given fresh and fresh water 3-4 times a day. Keep the animals in a clean place. To prevent foot and mouth disease in animals, vaccinate them with FMD vaccine and to prevent foot and mouth disease, vaccinate them with BQ vaccine. Make sure to give clean and fresh water to the animals 3-4 times a day.

Poultry Specific Advisory:

Poultry	Poultry Specific Advisory
HEN	Farmers are advised to the hen supplements the food, mix vitamins and energy food material in their feed as well as calcium content to hens. To prevent Ranikhet disease in chickens, vaccinate one week old chicks with F-1 vaccine and four week old chicks with R-2 vaccine.

Likely impacts of weather warnings (General)

According to the weather forecast received from the Indian Meteorological Department, there is no warning in next five days.

Impact based advisories (General)

Keeping in view the weather conditions, maintain proper moisture in summer vegetables and crops by doing light irrigation as per requirement. And harvest and thresh mature crops like mustard, gram, peas should be harvested in the morning/evening and do not go out of the house unnecessarily during the afternoon. While working in the field, tie a towel on the head and keep a water bottle with you.

Farmers are advised to download Unified Mausam and "Meghdoot" android application on mobile for Weather forecast and weather based Agromet Advisories and "Damini" android application for forecast of Thunderstorm and lightning.

Mausam MobileApp link: <https://play.google.com/store/apps/details/>

Meghdoot MobileApp link: https://play.google.com/store/apps/details

Damini MobileApp link : https://play.google.com/store/apps/details